

# Volatile compounds of ditax fruit (*Detarium senegalense* J.F. Gmel) from Senegal

Nafissatou Diop NDIAYE<sup>1</sup>, Marc LEBRUN<sup>2</sup>, Manuel DORNIER<sup>3\*</sup>

<sup>1</sup> Inst. Technol. Alim., Route Pères Maristes, BP 2765, Dakar Hann, Sénégal

<sup>2</sup> CIRAD, UMR95 QualiSud, TA B-95/16, 73 av. Jean François Breton, F-34398 Montpellier cedex 5, France

<sup>3</sup> Montpellier SupAgro, UMR95 QualiSud, Inst. Rég. Chaudes, 1101 av. Agropolis, BP 5098, F-34093 Montpellier cedex 5, France, dornier@cirad.fr

## Volatile compounds of ditax fruit (*Detarium senegalense* J.F. Gmel) from Senegal.

**Abstract – Introduction.** *Detarium senegalense* J.F. Gmel is a forest tree found in Senegal whose fruits are locally called ditax in Wolof. It is eaten fresh but it is widely used as nectar, which is one of the most popular beverages in Senegal. However, the chemical characterization of ditax pulp remains incomplete. This paper describes the volatile compounds of ditax to assess its organoleptic qualities. **Materials and methods.** Free volatile compounds of fresh ditax pulp were isolated by solvent-assisted flavor evaporation and analysis by GC-MS. **Results and discussion.** Among the 53 compounds tentatively identified, 49 are reported for the first time in this fruit. In total, 17 aldehydes, 11 aliphatic alcohols, 1 terpene alcohol, 7 free fatty acids, 3 unsaturated hydrocarbons, 1 terpene hydrocarbon, 7 sesquiterpene hydrocarbons, 1 phenol, 2 ketones, 2 esters and 1 organic acid compound were tentatively identified in ditax fresh pulp. The main volatiles identified in fresh ditax pulp were *trans*, *cis*-2,6-nonadienal (2.47 mg·kg<sup>-1</sup>), *cis*-2-heptenal (1.93 mg·kg<sup>-1</sup>), *trans*- $\alpha$ -bergamotene (1.11 mg·kg<sup>-1</sup>), bicyclo [2,2,0] hexane-1-carboxaldehyde (0.80 mg·kg<sup>-1</sup>), butyl octadecanoate (0.55 mg·kg<sup>-1</sup>) and *trans*-2-nonenal (0.47 mg·kg<sup>-1</sup> fresh pulp). **Conclusion.** Among the volatile compounds identified, aldehyde compounds were widely predominant. To assess the aromatic qualities of ditax pulp, the primary impact aromas should be determined by identifying the aroma-active compounds by GC-olfactometry.

## Senegal / *Detarium senegalense* / fruits / flavor / volatile compounds / solvent extraction

### Les composés volatils du fruit de ditax (*Detarium senegalense* J.F. Gmel) au Sénégal.

**Résumé – Introduction.** *Detarium senegalense* J.F. Gmel est un arbre forestier, trouvé au Sénégal, dont les fruits sont localement appelés ditax en Wolof. Le fruit est directement consommé mais il est largement utilisé en nectar, l'une des boissons les plus populaires au Sénégal. Cependant, la caractérisation chimique de la pulpe de ditax reste incomplète. Cette étude décrit les composés volatils de la pulpe de ditax afin d'évaluer sa qualité organoleptique. **Matériel et méthodes.** Les composés volatils libres de la pulpe de ditax fraîche ont été isolés par la technique d'évaporation assistée par solvant et analysés par GC-MS. **Résultats et discussion.** Parmi les 53 composés identifiés, 49 sont signalés pour la première fois dans ce fruit. Au total, 17 aldéhydes, 11 alcools aliphatiques, 1 alcool terpénique, 7 acides gras libres, 3 hydrocarbures insaturés, 1 hydrocarbure terpénique, 7 hydrocarbures sesquiterpéniques, 1 phénol, 2 cétones, 2 esters et 1 acide organique ont été identifiés dans la pulpe de ditax fraîche. Les principaux composés volatils identifiés dans la pulpe de ditax fraîche ont été le *trans*, *cis*-2,6-nonadiénal (2,47 mg·kg<sup>-1</sup>), le *cis*-2-hepténal (1,93 mg·kg<sup>-1</sup>), le *trans*- $\alpha$ -bergamotène (1,11 mg·kg<sup>-1</sup>), le bicyclo [2,2,0] hexane-1-carboxaldéhyde (0,80 mg·kg<sup>-1</sup>), l'octadecanoate de butyle (0,55 mg·kg<sup>-1</sup>) et le *trans*-2-nonénal (0,47 mg·kg<sup>-1</sup> de matière fraîche). **Conclusion.** Parmi les composés volatils identifiés, les aldéhydes sont les composés majoritaires. Pour évaluer la qualité aromatique de pulpe de ditax, l'identification des composés d'impacts notamment par GC-olfactométrie serait une perspective intéressante.

### Sénégal / *Detarium senegalense* / fruits / flaveur / composé volatil / extraction par solvant

\* Correspondence and reprints

Received 2 September 2013  
Accepted 11 September 2013

Fruits, 2014, vol. 69, p. 181–188  
© 2014 Cirad/EDP Sciences  
All rights reserved  
DOI: 10.1051/fruits/2014007  
[www.fruits-journal.org](http://www.fruits-journal.org)

RESUMEN ESPAÑOL, p. 188

## 1. Introduction

The *Detarium senegalense* tree from the Ceasalpiniaceae family, an arboreal fruit species, is still found wild in Senegal (West Africa), in the Sine-Saloum islands and in Casamance [1]. Reaching (15 to 40) m, it is also found in the wet dense forest borders, the coastal and septentrional regions, and the Sudan-Guinean zone. The *Detarium senegalense* fruit, locally called ditax in Wolof, is one of the most important forest fruit-bearing species with great economic importance. The fruit (*figure 1*) appears in the form of a drupe with a diameter from (3 to 8) cm and a mean weight of  $(49.2 \pm 11.2)$  g. The epicarp (13% of the fruit weight) is dark green and hard in unripe fruits, and light green, brown and brittle in ripe fruits. The pulp (34% of the fruit weight) is green and entangled in a high-fiber network firmly inserted on the stone (53% of the fruit weight), which contains one single dark brown seed [1–4]. The pulp is very rich in ascorbic acid [from (1.2 to 2.2)  $\text{g}\cdot 100\text{ g}^{-1}$  of fresh weight according to geographical area]; sucrose (18–20% dry weight) is the main component of the total sugars (23% dry weight) and pheophytin a ( $128\text{ mg}\cdot\text{kg}^{-1}$  fresh weight) is the major pigment of ditax pulp [5, 6]. The fruit is eaten fresh but it is widely used as nectar (fruit pulp with water and sugar), which is one of the most popular beverages in Senegal. However, the chemical characterization of ditax pulp remains incomplete. To the best of our knowledge, only Haddad makes reference to the aroma volatiles of ditax [3]. Therefore, our paper describes for the first time the volatile compounds of ditax to complete the chemical characterization data of this forest fruit and to assess its organoleptic qualities.

## 2. Materials and methods

### 2.1. Fruits

Fresh ripe fruits were bought at the market in September 2012, and after one week of storage at  $-18\text{ }^{\circ}\text{C}$ , pulps were extracted just before analysis.

### 2.2. Solvent-assisted flavor evaporation (SAFE) of ditax pulp volatiles

The solvent-assisted flavor evaporation (SAFE) technique was applied for careful extraction of the ditax pulp volatiles. Volatiles, preliminarily extracted in a solvent, were evaporated at low temperature in high vacuum conditions ( $10^{-3}$  mbar) and condensed with a liquid nitrogen trap.

Forty g of fresh ditax pulp and 50  $\mu\text{L}$  of internal standard solution (20  $\mu\text{L}\cdot 20\text{ mL}^{-1}$   $\text{H}_2\text{O}$ ) of nonan-4-ol were combined with 40 mL of distilled water and 80 mL of dichloromethane, and stirred for 30 min in an iced water bath under a nitrogen atmosphere. The supernatant was collected and the residue was reextracted with 80 mL of dichloromethane and 40 mL of distilled water for 30 min again to achieve complete transfer of the volatile compound from the pulp. Organic phases were combined, dried over anhydrous sodium sulfate and distilled at  $30\text{ }^{\circ}\text{C}$ . After complete distillation, the system was left for 1 h under vacuum. The distillate was then concentrated in a water heating bath at  $40\text{ }^{\circ}\text{C}$  using a Kuderna-Danish apparatus. Preliminary extractions were realized using a mixture of pentane-diethyl ether (50/50).

### 2.3. GC-MS analysis of safe extract

A tandem gas chromatograph 6890 / MSD 5973 / Gerstel Multipurpose Sample MPS-2 (Agilent Technologies, Palo Alto, USA) was used for volatile compound analysis. Injection (1  $\mu\text{L}$ ) was carried out in split 1/10 mode at  $250\text{ }^{\circ}\text{C}$  using a DB-WAX polar column (J&W,  $30\text{ m} \times 0.25\text{ mm} \times 0.25\text{ }\mu\text{m}$ ) and a DB-1ms nonpolar column (J&W,  $0.25\text{ mm} \times 30\text{ m} \times 0.25\text{ }\mu\text{m}$ ) (Agilent Technologies, Palo Alto, USA). Helium was used as the carrier gas at  $1\text{ mL}\cdot\text{min}^{-1}$ . Elution was realized with the following temperature program:  $3\text{ }^{\circ}\text{C}$  per min from  $40\text{ }^{\circ}\text{C}$  to  $170\text{ }^{\circ}\text{C}$ , then  $10\text{ }^{\circ}\text{C}$  per min up to  $240\text{ }^{\circ}\text{C}$  and held for 10 min. Mass spectra were recorded in EI+ mode at 70 eV within (15 to 300) Da. The analyzer and source temperatures were  $150\text{ }^{\circ}\text{C}$  and  $250\text{ }^{\circ}\text{C}$ , respectively.

## 2.4. Identification of volatile compounds

Data were analyzed with the MSD Chem revD1 acquisition and data retreatment software. Peak identification was done by comparing mass spectra with those of the NIST 2008 (National Institute of Standard Technology) database [7]. Injection of n-alkane series C<sub>8</sub>-C<sub>20</sub> (Sigma-Aldrich) was used for retention index (RI) calculations and their comparison with those found in the literature, and on the Flavornet<sup>1</sup> and Pherobase<sup>2</sup> websites.

## 2.5. Quantification of volatile compounds

Semi-quantitative determinations were carried out by adding nonan-4-ol (Sigma Aldrich) as an internal standard at a concentration of 20 µL:20 mL<sup>-1</sup> H<sub>2</sub>O. The ratio of the response factor was considered to be equal to 1, thus allowing semi-quantification in µg·kg<sup>-1</sup> fresh pulp using the equation:

$$m_i (\mu\text{g}\cdot\text{kg}^{-1} \text{ fresh pulp}) = 1000 \times \frac{A_i m_{IS}}{A_{IS} m_p}$$

with  $A_i$  = peak area of volatile compound  $i$ ;  $A_{IS}$  = peak area of nonan-4-ol;  $m_{IS}$  = quantity of nonan-4-ol;  $m_p$  = pulp quantity.

## 3. Results and discussion

The aroma compounds were separated on two columns of different polarity (DB-WAX and DB-1ms) (table D). Quantification on both the DB-WAX and DB-1ms columns, (11 769 and 10 200) µg·kg<sup>-1</sup>, were quite similar. This slight difference could be

<sup>1</sup> Acree T., Arn H., Flavornet and human odor space, Gas chromatography-olfactometry (GCO) of natural products, Datu Inc. 2004, www.flavornet.org.

<sup>2</sup> El-Sayed A.M., The Pherobase : Database of Pheromones and Semiochemicals, www.pherobase.com, 2003–2012, The Pherobase.



**Figure 1.** Fruit of *Detarium senegalense* J.F. Gmel: whole fruit, husked fruit (green pulp), fruit after pulp extraction (stone with fibrous network).

explained by a longer storage in the freezer (17 d at -18 °C) before analysis on the DB-1ms. Among the 53 compounds tentatively identified, forty-nine are reported for the first time in this fruit. In total, seventeen aldehydes, eleven aliphatic alcohols, one terpene alcohol, seven free fatty acids, three unsaturated hydrocarbons, one terpene hydrocarbon, seven sesquiterpene hydrocarbons, one phenol, two ketones, two esters and one organic acid compound were identified or tentatively identified in ditax fresh pulp (table D). Aldehydes known to impart “green” and “grassy” notes were qualitatively and quantitatively the most important class of compounds in ditax fresh pulp, with a total concentration of 6 704 µg·kg<sup>-1</sup> of fresh pulp (57% of the total volatile fraction). Among them, *trans*, *cis*-2,6-nonadienal was the main volatile, representing 21% of the total volatile fraction. Then followed *cis*-2-heptenal (16.4% of the total volatile fraction), bicyclo [2,2,0] hexane-1-carboxaldehyde (6.8%), *trans*-2-nonenal (4%) and *trans*-2-hexenal (3.8%).

**Table 1.** Volatiles compounds identified or tentatively (\*) in fresh ditax pulp after solvent-assisted flavour evaporation (SAFE) extraction and analysis on polar and non polar column.

| Compounds                                | Retention indice polar <sup>a</sup> |                         | Retention indice non polar <sup>b</sup> |                         | Concentration<br>( $\mu\text{g}\cdot\text{kg}^{-1}$ of fresh pulp) | Odor   |
|--|-------------------------------------|-------------------------|---|-------------------------|--|--|
|  | Experienced                         | Literature <sup>c</sup> | Experienced                             | Literature <sup>c</sup> |  |  |
| <b>Aliphatic alcohols</b>                |                                     |                         |   |                         |  |  |
| 1-Butanol                                | 1144                                | 1145                    | –                                       | 658                     | 12.34 ± 0.66   | Medicine, fruit <sup>1</sup> [8]             |
| 1-Penten-3-ol                            | 1155                                | 1181                    | –                                       | 669                     | 77.87 ± 2.76   | –  |
| 1-Pentanol                               | 1242                                | 1255                    | 814                                     | 730                     | 52.60 ± 3.75   | Balsamic <sup>1</sup>                        |
| Cis-2-penten-1-ol                        | nd                                  | 1313                    | 814                                     | 740                     | 112.12   | –  |
| Cyclopentanol                            | 1314                                | 1323                    | –                                       | 774                     | 159.99 ± 4.43  | –  |
| Cis-3-hexen-1-ol                         | 1384                                | 1401                    | 859                                     | 844                     | 67.50 ± 5.05   | Leaf, fatty grass [9]                        |
| 1-Hexanol                                | 1353                                | 1360                    | 870                                     | 858                     | 332.18 ± 10.98   | Fruit [9]; resin, flower, green <sup>1</sup> |
| Trans-2-hexen-1-ol                       | 1410                                | 1400                    | 867                                     | 870                     | 49.14 ± 0.34   | Fruit, leaf [9]; green [8]                   |
| 1-Heptanol                               | 1469                                | 1467                    | 950                                     | 945                     | 67.11 ± 0.37   | Chemical, green <sup>1</sup>                 |
| 6-Hepten-1-ol                            | 1530                                | –                       | 948                                     | 950                     | 35.82 ± 0.16   | –  |
| Cis-6-nonen-1-ol                         | 1756                                | 1714                    | 1151                                    | 1171                    | 73.64 ± 0.60   | –  |
| <b>Terpenic Alcohols</b>                 |                                     |                         |   |                         |  |  |
| 1,8-Cineole                              | 1194                                | 1214                    | 1009                                    | 1015                    | 31.85 ± 1.27   | Mint, sweet <sup>1</sup>                     |
| <b>Aldehydes</b>                         |                                     |                         |   |                         |  |  |
| Pentanal                                 | 937                                 | 935                     | –                                       | 674                     | 105.79 ± 6.94  | Almond, malt, pungent <sup>1</sup>           |
| Hexanal                                  | 1096                                | 1093                    | 824                                     | 773                     | 199.46 ± 9.40  | Green [8]; grass, tallow, fat <sup>1</sup> , |
| Trans-2-pentenal                         | 1126                                | 1117                    | 804                                     | 724                     | 18.08 ± 0.49   | Strawberry, fruit, tomato <sup>1</sup>       |
| Cis-3-hexenal                            | 1137                                | 1132                    | 823                                     | 801                     | 12.13 ± 0.14   | Leaf, green <sup>1</sup>                     |
| Heptanal                                 | 1172                                | 1174                    | 887                                     | 885                     | 72.97 ± 3.06   | Fat, citrus, rancid <sup>1</sup>             |
| Cis-2-hexenal                            | 1186                                | 1187                    | 847                                     | 815                     | Traces   | –  |
| Trans-2-hexenal                          | 1201                                | 1220                    | 851                                     | 826                     | 452.18 ± 19.58   | Green [8]                                    |
| Cis-4-heptenal                           | 1220                                | 1230                    | 878                                     | 885                     | 20.31 ± 0.79   | Biscuit, cream <sup>1</sup>                  |
| Cis-2-heptenal                           | 1310                                | 1319                    | 931                                     | 927                     | 1929.56 ± 21.96  | Green <sup>1</sup>                           |
| Bicyclo [2.2.0] hexane-1-carboxaldehyde* | 1372                                | –                       | 916                                     | –                       | 803.85 ± 1.68  | –  |
| Nonanal                                  | 1387                                | 1385                    | 1078                                    | 1079                    | 29.98 ± 0.11   | Fatty, citrus, green <sup>1</sup>            |
| Trans-2-octenal                          | 1425                                | 1442                    | 1024                                    | 1031                    | 16.00 ± 0.84   | –  |
| Cis, cis-3,6-nonadienal                  | nd                                  | 1444                    | 1065                                    | 1083                    | 34.17  | –  |
| Cis-6-nonenal                            | 1449                                | 1469                    | nd                                      | 1112                    | 27.34 ± 4.58   | –  |
| Trans, cis-2,4-heptadienal               | 1463                                | 1480                    | nd                                      | 921                     | 37.60 ± 1.08   | –  |
| Trans-2-nonenal                          | 1545                                | 1527                    | 1132                                    | 1130                    | 470.97 ± 1.96  | Cucumber, fatty, green <sup>1</sup>          |
| Trans, cis-2,6-nonadienal                | 1602                                | 1605                    | 1125                                    | 1137                    | 2474.08 ± 71.11  | Cucumber, waxy, green <sup>1</sup>           |

**Table I.**  
continued.

| Compounds                            | Retention indice polar <sup>a</sup> |                         | Retention indice nonpolar <sup>b</sup> |                         | Concentration<br>( $\mu\text{g}\cdot\text{kg}^{-1}$ of fresh pulp) | Odor                              |
|--------------------------------------|-------------------------------------|-------------------------|--|-------------------------|--|-----------------------------------|
|                                      | Experienced                         | Literature <sup>c</sup> | Experienced                            | Literature <sup>c</sup> |  |                                   |
| <b>Free fatty acids</b>              |                                     |                         |  |                         |  |                                   |
| Hexanoic acid                        | 1879                                | 1872                    | 1004                                   | 890                     | 273.24 $\pm$ 0.96  | Fat, sweat <sup>1</sup>           |
| Heptanoic acid                       | 1985                                | 1990                    | 1125                                   | 1064                    | 82.96 $\pm$ 4.24   | Rancid <sup>1</sup>               |
| <i>Trans</i> -2-hexanoic acid        | 1996                                | 1962                    | nd                                     | -                       | 20.02 $\pm$ 1.20   | -                                 |
| Nonanoic acid                        | 2183                                | 2202                    | 1280                                   | 1258                    | 26.73 $\pm$ 4.10   | Rancid <sup>1</sup>               |
| Dodecanoic acid*                     | 2613                                | 2517                    | 1501                                   | 1556                    | 33.17 $\pm$ 0.03   | Metal <sup>1</sup>                |
| Hexadecanoic acid                    | 2936                                | 2940                    | 1955                                   | 1950                    | 202.39 $\pm$ 13.24   | -                                 |
| Octadecanoic acid                    | 3184                                | 3181                    | 2093                                   | 2124                    | 200.59 $\pm$ 41.76   | -                                 |
| <b>Ketones</b>                       |                                     |                         |  |                         |  |                                   |
| 1-Penten-3-one                       | 1063                                | 1034                    | nd                                     | 673                     | 87.76 $\pm$ 2.99   | Fish, pungent <sup>1</sup>        |
| 3-Hydroxy-2-butanone                 | 1266                                | 1272                    | nd                                     | 678                     | 81.84 $\pm$ 5.97   | -                                 |
| <b>Esters</b>                        |                                     |                         |  |                         |  |                                   |
| Butyl hexadecanoate*                 | 2573                                | -                       | 2110                                   | 1977                    | 367.87 $\pm$ 3.11  | -                                 |
| Butyl octadecanoate*                 | 2733                                | -                       | 2257                                   | 2374                    | 552.07 $\pm$ 6.92  | -                                 |
| <b>Unsaturated hydrocarbons</b>      |                                     |                         |  |                         |  |                                   |
| Toluene                              | 1071                                | 1070                    | nd                                     | 756                     | 39.54 $\pm$ 1.62   | Paint <sup>1</sup>                |
| Ethylbenzene                         | 1121                                | 1124                    | 864                                    | 846                     | 5.63 $\pm$ 0.03  | -                                 |
| Styrene                              | 1234                                | 1261                    | 882                                    | 876                     | 154.47 $\pm$ 9.88  | Balsamic <sup>1</sup>             |
| <b>Terpene hydrocarbons</b>          |                                     |                         |  |                         |  |                                   |
| Limonene                             | 1182                                | 1178                    | 1012                                   | 1022                    | 5.5 $\pm$ 1  | Citrus, mint <sup>1</sup>         |
| <b>Sesquiterpene hydrocarbons</b>    |                                     |                         |  |                         |  |                                   |
| $\alpha$ -Copaene                    |                                     |                         |  |                         | 11.21  | Woody, spicy <sup>1</sup>         |
| <i>Cis</i> - $\alpha$ -bergamotene   | 1580                                | 1580                    | 1404                                   | 1414                    | 114.07 $\pm$ 3.06  | -                                 |
| $\alpha$ -Santalene                  | 1581                                | 1574                    | 1409                                   | 1419                    | 96.30 $\pm$ 6.15   | -                                 |
| <i>Trans</i> - $\alpha$ -bergamotene | 1602                                | 1779                    | 1425                                   | 1431                    | 1112 $\pm$ 6.17  | Woody, tea, warm <sup>1,2</sup>   |
| <i>Trans</i> - $\beta$ -farnesene*   | 1692                                | 1674                    | 1441                                   | 1448                    | 98.83 $\pm$ 7.75   | Woody, citrus, sweet <sup>1</sup> |
| $\beta$ -Bisabolene*                 | 1750                                | 1726                    | 1465                                   | 1513                    | 35.02 $\pm$ 4.31   | Balsamic <sup>1</sup>             |
| $\beta$ -Cadinene*                   | 1777                                | 1982                    | 1497                                   | 1440                    | 60.41 $\pm$ 0.94   | -                                 |
| <b>Phenol</b>                        |                                     |                         |  |                         |  |                                   |
| Phenol                               | 2026                                | 1479                    | 958                                    | 961                     | 43.52 $\pm$ 1.51   | Phenol <sup>1</sup>               |
| <b>Acid</b>                          |                                     |                         |  |                         |  |                                   |
| Acetic acid                          | 1450                                | 1452                    | nd                                     |                         |  | Acid <sup>1</sup>                 |

<sup>a</sup> linear retention index on DB-wax.

<sup>b</sup> linear retention index on DB-1ms.

<sup>c</sup> linear retention index from Flavornet<sup>1</sup>, Pherobase<sup>2</sup> and NIST 2008 [7].

trace: < 5  $\mu\text{g}\cdot\text{kg}^{-1}$  fresh pulp.

nd: Retention indice (RI) experimented < 800 or not detected.

Haddad reported that 2,6-nonadienal (42.3%), *trans*-2-heptanal (13.5%), *cis*-6-nonenal (7.5%), *trans*-2-nonenal (7.4%), caryophyllene oxide (2.5%), humulene oxide (1.8%) and tetradecane (0.7%) were the main volatiles of essential oil extract of ditax pulp obtained by hydrodistillation [3]. The following different flavors confirmed our perception; *trans*, *cis*-2,6-nonadienal is cucumber, violet, green and waxy; *trans*-2-nonenal is fatty, penetrating and waxy; melon or green melon and citrus are characteristic of *cis*-6-nonenal and green and watermelon-like of *cis*, *cis*-3,6-nonadienal [10–15]. Hexanal and *trans*-2-hexenal odor is green [8]. It has been widely reported that C<sub>9</sub> odorant compounds were the result of lipoxygenase degradation of linoleic and linolenic acid, which occurs rapidly after tissue disruption [16, 17].

Sesquiterpene hydrocarbons represent the second class of compounds (12.9% of total volatile compounds), with *trans*- $\alpha$ -bergamotene being the major component, with 73% of total sesquiterpene hydrocarbons and 9.4% of the total volatile fraction.

Aliphatic alcohols were the third class of compounds, with 8.8% of the total volatile fraction for a total amount of 1 040  $\mu\text{g}\cdot\text{kg}^{-1}$  of fresh pulp. Among them, 1-hexanol (floral, green notes) was the main volatile (2.8% of the total volatile fraction) followed by cyclopentanol (1.4%). However, other alcohols such as 1-butanol, *cis*-3-hexen-1-ol and *trans*-2-hexen-1-ol have also been identified in ditax pulp. They have already been reported in many other fruits such as apricot [18–21], copoazù [22] and acerola [23].

1,8-Cineole (mint, sweet flavor) was the only terpene alcohol identified in ditax pulp (0.3% of the total volatile fraction). 1,8-Cineole has also been identified in cantaloupe [24]. Alcohols and aldehydes of six carbon atoms are responsible for the herbaceous odor of several fruits and vegetables [20, 25].

Ester compounds were represented by butyl hexadecanoate (368  $\mu\text{g}\cdot\text{kg}^{-1}$ ) and butyl octadecanoate (552  $\mu\text{g}\cdot\text{kg}^{-1}$ ); these two compounds were tentatively identified because the experimental RIs (retention

indices) are different from those from the literature. These two compounds are esters of fatty acid and represent 7.8% of the total volatile fraction of ditax fresh pulp.

Free fatty acids represent 7% (839  $\mu\text{g}\cdot\text{kg}^{-1}$ ) of the total volatile fraction, where hexanoic acid (273  $\mu\text{g}\cdot\text{kg}^{-1}$ ), n-hexadecanoic acid (202  $\mu\text{g}\cdot\text{kg}^{-1}$ ) and octadecanoic acid (201  $\mu\text{g}\cdot\text{kg}^{-1}$ ) were the main compounds.

Acetic acid represents 2.45% of the total volatile fraction.

Other chemical compounds such as unsaturated hydrocarbons (1.7% of the total volatile fraction) with styrene as the main compound; ketones (1.45% of the total volatile fraction) and phenol (0.36% of the total volatile fraction) were also identified in ditax pulp.

## 4. Conclusion

The major volatiles present in ditax pulp could explain why ditax is very popular, especially for its strong cucumber-like aroma with gentle fruity, woody, herbal and green notes. Among the volatile compounds identified, aldehyde compounds were widely predominant. Five volatiles were responsible for over 38% of the total volatile fraction: *trans*, *cis*-2,6-nonadienal, (21%), *cis*-2-heptenal (16.4%), *trans*- $\alpha$ -bergamotene (9.4%) and bicyclo [2,2,0] hexane-1-carboxaldehyde (6.8%). To assess the aromatic qualities of ditax pulp, the primary impact aromas should be determined by identifying the aroma-active compounds by GC-olfactometry.

## Acknowledgments

The authors thank Mrs Oumar Diémé, Babacar Dieng and Noël Diandy from the Inst. Food Technol. (Dakar, Senegal) for their technical help and supplying fruits. We also thank CIRAD and WAAPP 2 for their financial and technical support.

## References

- [1] Diop N., Ndiaye A., Cisse M., Dieme O., Dornier M., Sock O., Le ditax (*Detarium senegalense* J. F. Gmel.): principales caractéristiques et utilisations au Sénégal, *Fruits* 65 (2010) 293–306.
- [2] Arbonnier M., Arbres, arbustes et lianes de zones sèches d'Afrique de l'Ouest, 2e éd., CIRAD–MNHN, Paris, France, 2002, 541 p.
- [3] Haddad C., *Fruitiers sauvages du Sénégal*, Univ. Montpellier I, Thèse, Montpellier, France, 2000, 372 p.
- [4] Kerharo J., Adam, J.G., *La pharmacopée sénégalaise traditionnelle : plantes médicinales et toxiques*, Ed. Vigot Frères, Paris, France, 1974, pp. 285–287.
- [5] Diop N., Dornier M., Dhuique-Mayer C., Prades A., Munier-Pantel S., Pélissier Y., Laroque M., Sock O., Caractérisation d'un fruit sauvage du Sénégal : le ditax (*Detarium senegalense* J.F. Gmel), in : Boëtsch G., Guerci A., Gueye L., Guisse A., *Les plantes du Sahel, usages et enjeux sociaux*, CNRS Ed., Paris, France, 2012, pp. 99–108.
- [6] Diop Ndiaye N., Dhuique-Mayer C., Cisse M., Dornier M., Identification and thermal degradation kinetics of chlorophyll pigments and ascorbic acid from ditax nectar (*Detarium senegalense* J.F. Gmel), *J. Agric. Food Chem.* 59 (22) (2011) 12018–12027.
- [7] Stein S., Mirokhin Y., Tchekhovskoi D., Mallard G., The NIST mass spectral search program for NIST/EPA/NIH mass spectral library, *Distrib. Agilent Technologies for use with the GC/MS and LC/MS Chemstation*, U.S.A., 2008.
- [8] Qiao Y., Xie B. J., Zhang Ya., Zhang Yu., Gang F., Yao X.L., Pan S.Y., Characterization of aroma active compounds in fruit juice and peel oil of Jincheng sweet orange fruit (*Citrus sinensis* (L.) Osbeck) by GC-MS and GC-O, *Molecules* 13 (2008) 1333–1344.
- [9] Ollé D., *Caractérisation des polysaccharides et des composés aromatiques de différents cultivars de mangue (*Mangifera indica* L.)*. Devenir de ces constituants lors de la préparation des concentrés aromatiques pulpeux, *Ecol. Ntl. Sup. Ind. Agric. Alim. (ENSIA)*, Thèse, Paris, France, 1997, 193 p.
- [10] Beaulieu J.C., Lea J.M., Characterization and semi quantitative analysis of volatiles in seedless watermelon varieties using solid-phase micro extraction, *J. Agric. Food Chem.* 54 (2006) 7789–7793.
- [11] Hwan Oh S., Seom Lim B., Jin Hong S., Koo Lee S., Aroma volatile changes of netted muskmelon (*Cucumis melo* L.) fruit during developmental stages, *Hortic. Environ. Biotechnol.* 52 (6) (2011) 590–595.
- [12] Yajima I., Sarakibara H., Ide J., Yanai T., Hayashi K., Volatile flavor components of watermelon (*Citrullus vulgaris*), *Agric. Biol. Chem. (Tokyo)* 49 (1985) 3145–3150.
- [13] Kim K.S., Lee H.J., Keem S.M., Volatile flavor components in watermelon (*Citrullus vulgaris* S.) and Oriental melon (*Cucumis melo* L.), *Korean J. Food Sci. Technol.* 31 (1999) 322–328.
- [14] Rymal K.S., Nakayama T.O.M.N., Identification of some volatile compounds from cucumber, *J. Agric. Food Chem.* 22 (4) (1974) 717–718.
- [15] Fross D.A., Dunstone E.A., Ramshaw E.H., Stark W., The flavor of cucumbers, *J. Food Sci.* 27 (1) (1962) 90–93.
- [16] Grosch W., Schwarz J., Linoleic and linolenic acid as precursors of the cucumber flavor, *Lipids* 6 (1971) 351–352.
- [17] Palma-Harris C., McFeeters R.F., Fleming H.P., Fresh cucumber flavor in refrigerated pickles: Comparison of sensory and instrumental analysis, *J. Agric. Food Chem.* 50 (2002) 4875–4877.
- [18] Solís-Solís H.M., Calderón-Santoyo M., Schorr-Galindo S., Luna-Solano G., Ragazzo-Sánchez J.A., Characterization of aroma potential of apricot varieties using different extraction techniques, *Food Chem.* 105 (2007) 829–837.
- [19] Takeoka G., Flath R., Mon T., Teranishi R., Guentert M., Volatile constituents of apricot (*Prunus armeniaca* L.), *J. Agric. Food Chem.* 38 (1990) 471–477.
- [20] Gómez E., Ledbetter C.A., Development of volatile compounds during fruit maturation: characterization of apricot and plum × apricot hybrids, *J. Sci. Food Agric.* 74 (1997) 541–546.
- [21] Guillot S., Peytavi L., Bureau S., Boulanger R., Lepoutre J.P., Crouzet J., Schorr-Galindo S., Aroma characterization of various apricot varieties using headspace-solid phase micro extraction combined with gas chromatography-mass spectrometry and gas chromatography-olfactometry, *Food Chem.* 96 (2006) 147–155.

- [22] Quijano C.E., Pino J.A., Volatile compounds of copoazú (*Theobroma grandiflorum* Schumann) fruit, *Food Chem.* 104 (2007) 1123–1126.
- [23] Boulanger R., Crouzet J., Identification of the aroma components of acerola (*Malpighia glabra* L.), *Food Chem.* 74 (2001) 209–216.
- [24] Beaulieu J.C., Grimm C.C., Identification of volatiles compounds in Cantaloupe at various developmental stages using solid phase micro extraction, *J. Agric. Food Chem.* 49 (2001) 1345–1352.
- [25] López-Tamames E., Carro-Mariño N., Ziya-Gunata Y., Sapis C., Baumes R., Bayonove C., Potential aroma in several varieties of Spanish grape, *J. Agric. Food Chem.* 45 (1997) 1729–1735.

### Los compuestos volátiles del fruto de ditax (*Detarium senegalense* J.F. Gmel) en Senegal.

**Resumen – Introducción.** *Detarium senegalense* J.F. Gmel es una especie forestal que se encuentra en Senegal, cuyos frutos son denominados localmente ditax en Wólof. El fruto se consume directamente, pero también se emplea ampliamente en forma de néctar, una de las bebidas más populares de Senegal. Sin embargo la caracterización química de la pulpa de ditax permanece incompleta. El presente trabajo describe los compuestos volátiles de la pulpa de ditax con el fin de evaluar su calidad organoléptica. **Material y métodos.** Los compuestos volátiles libres de la pulpa de ditax fresca se aislaron mediante la técnica de evaporación asistida con solvente y se analizaron mediante GC-MS. **Resultados y discusión.** Entre los 53 compuestos identificados, se señalaron 49 por primera vez en este fruto. En total se identificaron 17 aldehídos, 11 alcoholes alifáticos, 1 alcohol terpénico, 7 ácidos grasos libres, 3 hidrocarburos sesquiterpénicos, 1 fenol, 2 cetonas, 2 éster y 1 ácido orgánico en la pulpa fresca de ditax. Los principales volátiles identificados en la pulpa de ditax fresca fueron el *trans*, *cis*-2,6-nonadienal ( $2.47 \text{ mg}\cdot\text{kg}^{-1}$ ), el *cis*-2-heptenal ( $1.93 \text{ mg}\cdot\text{kg}^{-1}$ ), el *trans*- $\alpha$ -bergamoteno ( $1.11 \text{ mg}\cdot\text{kg}^{-1}$ ), el biciclo [2,2,0] hexano-1-carboxaldehído ( $0.80 \text{ mg}\cdot\text{kg}^{-1}$ ), el octadecanoato de butilo ( $0.55 \text{ mg}\cdot\text{kg}^{-1}$ ) y el *trans*-2-nonenal ( $0.47 \text{ mg}\cdot\text{kg}^{-1}$  de materia fresca). **Conclusión.** Entre los compuestos volátiles identificados, los aldehídos son los compuestos mayoritarios. Para evaluar la calidad aromática de pulpa de ditax, una perspectiva interesante sería la identificación de los compuestos de impactos, sobre todo por GC-olfactometría.

**Senegal / *Detarium senegalense* / frutas / sabor / compuesto volátil / extracción por disolventes**